



Ultrasound-guided FNA efficiency in diagnosing of metastatic nodes of oral squamous cell carcinoma: A review

Sona Rafieyan ¹, Yousef Kananizadeh ², Elahe Reyhani ^{3*}

1. Department of Oral and Maxillofacial Pathology, School of Dentistry, Zanjan University of Medical Sciences, Zanjan, Iran.

2. Department of Oral and Maxillofacial Surgery, Urmia University of Medical Sciences, Urmia, Iran.

3. Faculty of Dentistry, Zanjan University of Medical Sciences, Zanjan, Iran.

ARTICLE INFO

Article Type:
Review Article

Received: 2 Oct. 2020

Revised: 2 Dec. 2020

Accepted: 25 Mar. 2021

*Corresponding author:

Elahe Reyhani

Faculty of Dentistry, Zanjan University of Medical Sciences, Zanjan, Iran.

Tel: +98-21-84902473

Fax: +98-21-84902473

Email: s.rafieyan@Zums.ac.ir

ABSTRACT

Oral cancer comprises a group of neoplasms affecting any fields of the oral cavity, pharyngeal regions and salivary glands. However, this term tends to be used interchangeably with oral squamous cell carcinoma (OSCC), which represents the most common of all oral neoplasms. It is estimated that more than 90% of all oral neoplasms are OSCC. Oral SCC prefers to spread through the lymphatic system than the bloodstream, and in this region tongue tumors have the greatest potential for neck metastases. Primary tumor's histopathologic features like depth of invasion, thickness, growth pattern, differentiation, lymphovascular or vascular invasion have a limited prognostic value for nodal disease. Cervical lymph node status accurate assessment in HNSCC patients is critical issue because of its influences on prognosis and treatment decisions, USgFNAC proved widely accepted not only because of safety and availability in diagnosis but also it is repeatable, minimally invasive and cost-effectiveness.

Keywords: Squamous cell carcinoma; Hhead and neck cancer; Ultrasound-guided fine needle aspiration cytology; Ultrasound; Sentinel lymph node biopsy.

Introduction

Oral cancer comprises a group of neoplasms affecting any fields of the oral cavity, pharyngeal regions and salivary glands. However, this term tends to be used interchangeably with oral squamous cell carcinoma (OSCC), which represents the most frequent of all oral neoplasms. It is estimated that more of 90% of all oral neoplasms are OSCC [1]. It usually occurs after the

fifth decade of life, and is more common in men. They account for 5% of all malignancies worldwide and 3% in the United States [2,3]. Some of the proven main risk factors are: tobacco and alcohol consumption, Epstein-Barr virus (EBV) and human papilloma virus (especially 16 and 18) infections, and reverse smoking [4]. Current recommended treatment strategy is surgical resection with or without

chemotherapy and radiation therapy. The 5-year survival rate for oral and pharyngeal cancers is 56%, whereas it is 62% in laryngeal cancers. Two third of these patients present debilitating symptoms that impact their quality of life [5]. The oral cavity is full of lymphatic drainage, and regional nodal metastasis occurs early and constantly in oral SCC [6]. Cervical lymph node status accurate assessment in HNSCC patients is critical issue because of its influences on prognosis and treatment decisions [7]. Oral SCC prefers to spread through the lymphatic system than hematogenously. In this region tongue tumors have the greatest potential for neck metastases [8,9]. Early diagnosis of neck metastasis is an important prognostic factor. The survival rate of patients with Single ipsilateral or contralateral metastatic node decreases by 50% and it further reduces by 50% if there are bilateral metastatic nodes [10]. Surgical excision is essential to control primary tumor in SCC, recurrence often happens in cervical lymph nodes, which is the most important factor in the long-term prognosis of patients [11,12]. Primary tumor's histopathological features like depth of invasion, thickness, growth pattern, differentiation, lymphovascular or vascular invasion, extent of inflammatory reaction around the tumor, perineural invasion and numbers of biomarkers have a limited prognostic value for nodal disease [13]. According to latest articles, neck dissection (ND) in oral cancer patients with nodal involvement is the most suggested treatment method, but there is not an agreement to treat all patients with a clinically negative neck (cN0) with ND.

Lymph nodes excisional biopsy in cNo patients should be avoided because of potential of local complications (like hematoma, infection, disruption of fascial planes for subsequent neck dissection), as well as the theoretical risk of neck recurrence [14] Using selective neck dissection as a treatment could have risk of morbidity [15]. For example "shoulder syndrome" has been reported as a iatrogenic injury that occurs in spinal accessory nerve injury [16]. Carlos Moreno-García in 2016 reported 23.7-42% of recurrence in N0 neck [13]. If the "wait and see" policy is provided for cN0 patients occult neck metastasis can occur. The site, stage, and other characteristics of tumor are the risk factors of occult metastasis [15]. By following-up cN0 patients closely with the help of diagnostic techniques, such as ultrasound-guided fine-needle aspiration cytology (USgFNAC), occult metastases can be detected at early stages [16]. Screening of clinically N0 neck with ultrasound, magnetic resonance imaging (MRI), positron emission tomography (PET) or computed tomography

(CT) could be helpful in detection of some non-palpable nodal metastases, but their Limited accuracy compromise applying them. According to P Dabirmoghaddam et al in 2014 USgFNAC has a higher sensitivity and specificity to palpation and US alone and has an accuracy of 96%. In palpable metastatic lymph nodes, the sensitivity of USgFNA is often reported from 98% to 100% and 70% in nonpalpable lymph nodes. Adding cytology method increase diagnosis of malignant tumor specificity up to 100%. However, in patients with previous irradiation of the neck false positive findings of FNA have been reported [17]. A meta-analysis assessing all above-mentioned modalities (with the exception of PET CT scan) reported that USgFNAC has the highest diagnostic odds ratio for staging of neck with a pooled sensitivity of 80% and specificity of 98% for it [18]. The other meta-analysis reported PET-CT with few accuracy (specificity & sensitivity) in staging the cN0 patients [19]. Many clinicians still believe in surgery procedure in patient with overt lymph node metastases with or without adjuvant chemotherapy and radiation, But the cost-benefit of this treatment for cN0 patients should be considered [20]. The purpose of this review is evaluating the feasibility and performance of USgFNAC in Diagnosing of Metastatic Nodes of Oral Squamous Cell Carcinoma.

Discussion

-SLNB

The first lymph node in a drain of metastatic tumor cells is the sentinel lymph node, biopsy of this node is an early detective technique to identify the presence of nodal metastasis. Sentinel lymph node biopsy (SLNB) for oral SCC is not yet clearly defined because of the complex nature of head and neck region like various draining lymphatic pathways and high false-positive or negative rate of reports. Studies found SLNB as a valid diagnostic technique that properly stages regional metastases in HNSCC [21,22]. Although operative morbidity is potentially high, SLNB reported as an accurate staging procedure for cN0 patients [23]. A research with huge statically population reported the sensitivity of 93% in SLNB for early stage (T1-T2) oral and oropharyngeal cancers [24]. Also a meta-analysis of 21 studies concluded a 92% sensitivity for SLNB for staging the neck in oral cavity SCC [25]. The SLNB specificity of major studies is over 90%. Chaturvedi et al reported the sensitivity of SLNB (71%) that sound lower in comparison to USgFNAC [25,26]. The results of the two meta-analyses of Bondt et al suggest that efficacy of SLNB and USgFNAC is comparable [18,25].

Although SLNB has high accuracy, but it is not use commonly in most centers because of its time consuming and invasive procedure which requires multidisciplinary expertise [27].

-Ultrasonography (US)

Ultrasonography (US) is a simple method used for detection of noninvasive and soft tissue-related diseases in oral and maxillofacial regions [28]. The ultrasound image reveals the surface structures of computed tomography (CT) and magnetic resonance (MR). US may be clinically applied to assess lymph node-related diseases, salivary gland-related diseases, subcutaneous diseases, and tongue-related diseases [29]. Ultrasound of the neck has been used extensively for cervical metastases in head and neck cancer [30]. Ultrasound has higher sensitivity (96.8%) than palpation (73.3%) for detection of cervical lymph nodes [31]. De Bondt et al meta-analytic study in 2007 reported sensitivity and specificity of 66% and 78% for US. they Also reported that US is the most sensitive imaging technique for neck node staging with a pooled sensitivity of 87% and pooled specificity of 86% respectively [18,19]. according to results of some other research, a higher sensitivity of US has been reported in occult metastasis detection in the clinically negative neck [32-34]. Byers et al reported a sensitivity of 27% and specificity of 71% for the Ultrasound in the clinically negative neck, while To E et al reported a sensitivity of 47%. The result of both last researches declares limited prediction of Ultrasound in evaluating occult metastasis in tongue cancer [35,36].

Fine-needle aspiration (FNA) is a diagnostic method used to investigates masses. Fine-needle aspiration biopsies are very safe, less traumatic and minor surgical procedure with a few complications include bruising and soreness. Often excisional biopsy can be avoided by performing a needle aspiration biopsy instead, without hospitalization. FNA avoids the chance of seeding cancer cells in neck and making following treatment of a confirmed malignancy more challenging [35]. In some respects because of small size of sample problematic cells might be missed which could have false negative results [37]. However, FNAC has high efficacy and relative safety and comparing to open biopsy, this is comparatively inexpensive, well tolerated by patients with few complications without needs for general anesthesia [20,38]. In last decade fine needle aspiration cytology has been approved as a gold standard for diagnosing palpable head and neck lesions in the first step [13,39]. Using ultrasound with fine needle aspiration cytology offers the benefit of cytologic

analysis of suspicious lymph nodes and the presence of cancer cells and atypical cells in the pathology result of Ultrasound-guided fine-needle aspiration cytology (USgFNAC) was regarded as a positive result; otherwise, the result was considered negative [36]. Which has improved the specificity and overall accuracy of ultrasonography. Lymph nodes cytological analysis has an extremely high specificity that could approve lymph node metastasis presence in patients with HNSCC [19]. Whereas from the cellular material obtained through USgFNAC, different molecular examination like HPV analysis and tumor DNA sequencing can be achieved [40].

Diagnosing methods like Computed tomography (CT), magnetic resonance imaging (MRI), and positron emission tomography (PET) increase the detection certainty of involved lymph nodes [41,42]. In evaluation of nodal metastasis Ultrasound Guided FNAC is comparable to CT scan and can provide further information about enlarged lymph nodes and can show malignancy in small lymph nodes not found by CT or MRI. Also, clinician can present treatment plan to the cN0 patients before they leave the clinic, which avoids delay [43]. USgFNAC is a rapid and inexpensive procedure without the need of any contrast medium injection [44,45]. Some studies have reported low sensitivity of USgFNAC in clinically negative necks.

The confirmation of high accuracy of USgFNAC was based on 3 meta-analytic studies, 2 of them had included both clinically positive and negative necks [34,46]. It is clear that the larger the size of nodes with metastasis involvement, the more sensitive the USgFNAC technique gets. The third study of this meta-analysis was restricted to USgFNAC in cN0 patients and it reported a lower sensitivity of 48% and specificity of 100%, and overall accuracy of 79% [47]. Some studies also reported low sensitivity of USgFNAC in the estimate of cN0 patients [48,49]. Employing imaging in the neck nodes depends on some factors like the location and clinical extent of the primary tumor. Moreover, costing is also an important factor. USgFNAC can be recommended as a low-priced and most achievable added to the CT scan in the evaluation of neck node metastasis. Currently, the most reliable principle for assessing the nature of a lymph node and lymph node selection for aspiration is its size, measured as the minimal axial diameter. Based on studies, in order to obtain ideal sensitivity lymph nodes with a minimal axial diameter larger than 4mm in level II, and 3mm in other levels of the neck should be selected for ultrasound-guided FNAC. Selection criteria in the head

and neck region can estimate the overall accuracy of USgFNAC [50]. Ultrasound-guided FNAC decreases the risk of occult lymph node metastasis to 21% in the early oral squamous cell carcinoma [27,51]. Provides a plan for surgical intervention in malignant lesions and prevents unnecessary immediate surgery in benign nodules [52]. The limitation of USgFNAC is inaccessibility of paratracheal, retropharyngeal, and mediastinal lymph nodes that cause the main drawbacks of this technique [44,51]. Ultrasound-guided FNAC sensitivity has been reported as high as 73% in patients with a clinically (palpably) negative neck in head and neck squamous cell carcinoma [48]. Most studies identified USgFNA an appropriate technique because of the early availability of results, effortlessness, least trauma or injury, the low rate of complications and not needing heavy sedation or general anesthesia. USgFNA of the head and neck region is a well-accepted technique that has high specificity [53].

Conclusion

USgFNAC is an acceptable method that will be preferred in neck masses diagnosis especially in malignant lesions and it is widely accepted not only because of safety and availability in diagnosis but also it is repeatable, minimally invasive, non-ionizing and cost-effective [44,45]. Furthermore, USgFNAC is a reliable diagnosing method that enhances clinical researchers to collaborate with surgeons and also open the knowledge boundaries to optimistic studies on modifying the diagnosing method for earlier and exact treatment for SCC.

Conflict of Interest

There is no conflict of interest to declare.

References

[1] Markopoulos AK. Current aspects on oral squamous cell carcinoma. *The open dentistry journal*. 2012; 6:126.

[2] Siegel R, Ma J, Zou Z, Jemal AJ. *Cancer statistics, 2014*. 2014; 64(1):9-29.

[3] Ferlay J, Shin HR, Bray F, Forman D, Mathers C, Parkin DM. *Estimates of worldwide burden of cancer in 2008: GLOBOCAN 2008*. 2010; 127(12):2893-917.

[4] Chua ML, Wee JT, Hui EP, Chan AT. *Nasopharyngeal carcinoma*. 2016; 387(10022):1012-24.

[5] Reyes-Gibby CC, Anderson KO, Merriman KW,

Todd KH, Shete SS, Hanna EY. *Survival patterns in squamous cell carcinoma of the head and neck: pain as an independent prognostic factor for survival*. 2014; 15(10):1015-22.

[6] Massey C, Dharmarajan A, Bannuru RR, Rebeiz E. *Management of N0 neck in early oral squamous cell carcinoma: A systematic review and meta-analysis*. *The Laryngoscope*. 2019; 129(8):E284-E98.

[7] Peltenburg B, de Keizer B, Dankbaar JW, de Boer M, Willems SM, Philippens ME, et al. *Prediction of ultrasound guided fine needle aspiration cytology results by FDG PET-CT for lymph node metastases in head and neck squamous cell carcinoma patients*. *Acta Oncologica*. 2018; 57(12):1687-92.

[8] Montero PH, Patel SG. *Cancer of the oral cavity*. 2015; 24(3):491-508.

[9] Rodrigo JP, Grilli G, Shah JP, Medina JE, Robbins KT, Takes RP, et al. *Selective neck dissection in surgically treated head and neck squamous cell carcinoma patients with a clinically positive neck: Systematic review*. *European Journal of Surgical Oncology*. 2018; 44(4):395-403.

[10] Gubhaju M, Bajracharya A. *The role of ultrasound and ultrasound guided fine needle aspiration cytology in detection and management of neck node metastases in clinically N0 patients with carcinoma of upper aero-digestive tract in remote part of underdeveloped country*. *Annals of Oncology*. 2018; 29(suppl_9):mdy438. 025.

[11] Johnson JT, Barnes EL, Myers EN, Schramm VL, Borochoviz D, Sigler BA. *The extracapsular spread of tumors in cervical node metastasis*. *Archives of Otolaryngology*. 1981; 107(12):725-9.

[12] Aygun C, Salazar OM, Sewchand W, Amornmarn R, Prempre T. *Carcinoma of the floor of the mouth: a 20-year experience*. *International Journal of Radiation Oncology* Biology* Physics*. 1984; 10(5):619-26.

[13] Moreno-García C. *Elective neck dissection in early oral squamous cell carcinoma: Necessary*. 2016; 3:167-74.

[14] Golusinski P, Di Maio P, Pehlivan B, Colley S, Nankivell P, Kong A, et al. *Evidence for the approach to the diagnostic evaluation of squamous cell carcinoma occult primary tumors of the head*

- and neck. *Oral oncology*. 2019; 88:145-52.
- [15] Bradley PJ, Ferlito A, Silver CE, Takes RP, Woolgar JA, Strojan P, et al. Neck treatment and shoulder morbidity: still a challenge. 2011; 33(7):1060-7.
- [16] Popovski V, Benedetti A, Popovic-Monevska D, Grcev A, Stamoski A, Zhivadnovik J. Spinal accessory nerve preservation in modified neck dissections: surgical and functional outcomes. *Acta Otorhinolaryngologica Italica*. 2017; 37(5):368.
- [17] Dabirmoghaddam P, Sharifkashany S, Mashali LJIJoR. Ultrasound-guided fine needle aspiration cytology in the assessment of cervical metastasis in patients undergoing elective neck dissection. 2014; 11(3).
- [18] De Bondt R, Nelemans P, Hofman P, Casselman J, Kremer B, van Engelshoven J, et al. Detection of lymph node metastases in head and neck cancer: a meta-analysis comparing US, USgFNAC, CT and MR imaging. 2007; 64(2):266-72.
- [19] Liao L-J, Lo W-C, Hsu W-L, Wang C-T, Lai M-SJBc. Detection of cervical lymph node metastasis in head and neck cancer patients with clinically N0 neck—a meta-analysis comparing different imaging modalities. 2012; 12(1):236.
- [20] Huang SHJMo, *patologia oral y cirugia bucal*. Oral cancer: Current role of radiotherapy and chemotherapy. 2013; 18(2):e233.
- [21] Thompson C, John MS, Lawson G, Grogan T, Elashoff D, Mendelsohn A. Diagnostic value of sentinel lymph node biopsy in head and neck cancer: a meta-analysis. *European Archives of Oto-Rhino-Laryngology*. 2013; 270(7):2115-22.
- [22] Mehta V, Nathan CA. What is the role of sentinel lymph node biopsy in early-stage oral cavity carcinoma? *Laryngoscope*. 2016; 126(1):9-10.
- [23] Liu YY, Rozen WM, Rahdon RJ Ar. Sentinel lymph node biopsy for squamous cell carcinoma of the extremities: case report and review of the literature. 2011; 31(4):1443-6.
- [24] Ross GL, Soutar DS, MacDonald DG, Shoaib T, Camilleri I, Robertson AG, et al. Sentinel node biopsy in head and neck cancer: preliminary results of a multicenter trial. 2004; 11(7):690-6.
- [25] Govers TM, Hannink G, Merckx MA, Takes RP, Rovers MMJOO. Sentinel node biopsy for squamous cell carcinoma of the oral cavity and oropharynx: a diagnostic meta-analysis. 2013; 49(8):726-32.
- [26] Chaturvedi P, Datta S, Arya S, Rangarajan V, Kane SV, Nair D, et al. Prospective study of ultrasound-guided fine-needle aspiration cytology and sentinel node biopsy in the staging of clinically negative T1 and T2 oral cancer. 2015; 37(10):1504-8.
- [27] Amersi F, Morton DLJAis. The role of sentinel lymph node biopsy in the management of melanoma. 2007; 41:241-56.
- [28] Mashkevich G, Wang J, Rawnsley J, Keller GS. The utility of ultrasound in the evaluation of submental fullness in aging necks. *Archives of facial plastic surgery*. 2009; 11(4):240-5.
- [29] McKenna J, Friesen JR, Sowa MG, Hayakawa T, Kerr PD, Pabbies A. Assessing flap perfusion: optical spectroscopy versus venous doppler ultrasonography. *Journal of Otolaryngology--Head & Neck Surgery*. 2009; 38(5).
- [30] Wakasugi-Sato N, Kodama M, Matsuo K, Yamamoto N, Oda M, Ishikawa A, et al. Advanced clinical usefulness of ultrasonography for diseases in oral and maxillofacial regions. *International journal of dentistry*. 2010; 2010.
- [31] Metha SS, Mhapuskar AA, Marathe SP, Metha SS, Jadhav S, Thakare SD, et al. Assessment of efficacy of ultrasonography in cervical lymphadenopathy in oral malignancies. *Journal of family medicine and primary care*. 2019; 8(2):544.
- [32] Hodder S, Evans R, Patton D, Silvester KJBjoo, *surgery m*. Ultrasound and fine needle aspiration cytology in the staging of neck lymph nodes in oral squamous cell carcinoma. 2000; 38(5):430-6.
- [33] Thomsen JB, Sørensen JA, Grupe P, Karstoft J, Krogdahl AJAR. Staging N0 oral cancer: lymphoscintigraphy and conventional imaging. 2005; 46(5):492-6.
- [34] Knappe M, Louw M, Gregor RTJAoOH, *Surgery N*. Ultrasonography-guided fine-needle aspiration for the assessment of cervical metastases. 2000; 126(9):1091-6.
- [35] Akoğlu E, Dutipek M, Bekiş R, Değirmenci B, Ada E, Güneri AJJoo. Assessment of cervical lymph

- node metastasis with different imaging methods in patients with head and neck squamous cell carcinoma. 2005; 34(6).
- [36] Eskander A, Monteiro E, O'Connell D, Taylor SM. Head and Neck Surgical Oncology Choosing Wisely Campaign: imaging for patients with hoarseness, fine needle aspiration for neck mass, and ultrasound for odynophagia. *Journal of Otolaryngology-Head & Neck Surgery*. 2018; 47(1):2.
- [37] Dean DS, Gharib H. Fine-needle aspiration biopsy of the thyroid gland. *Endotext* [Internet]: MD-Text. com, Inc.; 2015.
- [38] Akkina SR, Kim RY, Stucken CL, Pynnonen MA, Bradford CR. The current practice of open neck mass biopsy in the diagnosis of head and neck cancer: A retrospective cohort study. *Laryngoscope Investigative Otolaryngology*. 2019; 4(1):57-61.
- [39] Haynes J, Arnold KR, Aguirre-Oskins C, Chandra S. Evaluation of neck masses in adults. *American family physician*. 2015; 91(10).
- [40] Rizzo G, Black M, Mymryk J, Barrett J, Nichols A. Defining the genomic landscape of head and neck cancers through next-generation sequencing. *Oral diseases*. 2015; 21(1):e11-e24.
- [41] Jiang P, Wang J, Ran W, Jiang Y, Tian S, Sun H. Five-year outcome of ultrasound-guided interstitial permanent 125I seeds implantation for local head and neck recurrent tumors: a single center retrospective study. *Journal of contemporary brachytherapy*. 2019; 11(1):28.
- [42] Brouwer J, de Bree R, Comans EF, Castelijns JA, Hoekstra OS, Leemans CR. Positron emission tomography using [18 F] fluorodeoxyglucose (FDG-PET) in the clinically negative neck: is it likely to be superior? *European Archives of Oto-Rhino-Laryngology and Head & Neck*. 2004; 261(9):479-83.
- [43] Schmidt RL, Jedrzkiwicz JD, Allred RJ, Matsuo S, Witt BLJH, neck. Verification bias in diagnostic accuracy studies for fine-and core needle biopsy of salivary gland lesions in otolaryngology journals: A systematic review and analysis. 2014; 36(11):1654-61.
- [44] Mangal N, Sharma VK, Verma N, Agarwal AK, Sharma SP, Aneja SJJ, O'Connell D. Ultrasound guided fine needle aspiration cytology in the diagnosis of retroperitoneal masses: A study of 85 cases. 2009; 26(3):97.
- [45] Ahuja AT, Ying M. Sonographic evaluation of cervical lymph nodes. *American Journal of Roentgenology*. 2005; 184(5):1691-9.
- [46] Van den Brekel M, Castelijns J, Stel H, Golding R, Meyer C, Snow GJE, et al. Modern imaging techniques and ultrasound-guided aspiration cytology for the assessment of neck node metastases: a prospective comparative study. 1993; 250(1):11-7.
- [47] Takes RP, Righi P, Meeuwis CA, Manni JJ, Paul Knegt M, Marres HA, et al. The value of ultrasound with ultrasound-guided fine-needle aspiration biopsy compared to computed tomography in the detection of regional metastases in the clinically negative neck. 1998; 40(5):1027-32.
- [48] Borgemeester MC, van den Brekel MW, van Tinteren H, Smeele LE, Pameijer FA, van Velthuysen MLF, et al. Ultrasound-guided aspiration cytology for the assessment of the clinically N0 neck: Factors influencing its accuracy. 2008; 30(11):1505-13.
- [49] Righi PD, Kopecky KK, Caldemeyer KS, Ball VA, Weisberger EC, Radpour SJH, et al. Comparison of ultrasound-fine needle aspiration and computed tomography in patients undergoing elective neck dissection. 1997; 19(7):604-10.
- [50] Silas O, Ige O, Adoga A, Nimkur L, Ajetunmobi OJJ, et al. Role of fine needle aspiration cytology (FNAC) as a diagnostic tool in paediatric head and neck lymphadenopathy. 2015; 4(1).
- [51] Krane JF. Role of cytology in the diagnosis and management of HPV-associated head and neck carcinoma. *Acta cytologica*. 2013; 57(2):117-26.
- [52] Gupta A, Ly S, Castroneves LA, Frates MC, Benson CB, Feldman HA, et al. A standardized assessment of thyroid nodules in children confirms higher cancer prevalence than in adults. *The Journal of Clinical Endocrinology & Metabolism*. 2013; 98(8):3238-45.
- [53] Sergi C, Dhiman A, Gray J-AJD. Fine Needle Aspiration Cytology for Neck Masses in Childhood.

An Illustrative Approach. 2018; 8(2):28.

Please cite this paper as:

Rafieyan S, Kananizadeh Y, Reyhani E; Ultrasound-guided FNA efficiency in diagnosing of metastatic nodes of oral squamous cell carcinoma: A review. *J Craniomaxillofac Res* 2021; 8(2): 44-50