



The Role of Platelet-Rich Products in Soft Tissue Repair after Extraction of Wisdom Teeth: A Review

Farzaneh Bolandparva^{1,2}, Nima Malekhosseini^{3*} , Arvin Asgarzade³

1. Department Oral and Maxillofacial Surgery, School of Dentistry, Tehran University of Medical Sciences, Tehran, Iran.

2. Craniomaxillofacial Research Center, Tehran University of Medical Sciences, Tehran, Iran.

3. DDs, School of Dentistry, Tehran University of Medical Sciences, Tehran, Iran.

ARTICLE INFO

Article Type:
Review Article

Received: 12 October 2024

Revised: 12 November 2024

Accepted: 19 December 2024

**Corresponding author:*

Nima Malekhosseini

Department of Oral and Maxillofacial Surgery,
School of Dentistry, Tehran University of Medical
Sciences, Tehran, Iran.

Tel: +98-912-0993286

Fax: +98-21-84903747

Email: greennima2@gmail.com

ABSTRACT

Introduction: The extraction of the third molar, commonly known as wisdom tooth extraction, is one of the most frequent clinical procedures. This surgery can lead to several postoperative complications such as pain, trismus, swelling, dry socket and damage to soft and hard tissues, significantly impacting the patient's quality of life. Platelet-Rich Plasma (PRP) and Platelet-Rich Fibrin (PRF) have been proposed as an effective treatment to mitigate these complications and enhance tissue repair. This study aims to review the role of PRP and PRF in soft tissue repair following wisdom tooth extraction between 2015 and 2023.

Materials and Methods: This research reviewed the role of PRP and PRF in soft tissue repair after wisdom tooth extraction. Articles from Scopus, PubMed, Elsevier, and SID databases were selected and reviewed based on inclusion and exclusion criteria, focusing on publications from 2015 to 2023. The search was conducted using keywords such as "impacted wisdom teeth," "soft tissue," "hard tissue," "repair," "PRP" and "PRF."

Results: Nineteen relevant studies were identified, with no studies conducted in 2023. The studies from this period were primarily meta-analyses with varying degrees of relevance to the study topic. The distribution of the studies reviewed is as follows: three studies in 2022, four studies in 2021, three studies in 2020, one study in 2019, one study in 2018, one study in 2017, two studies in 2016, and four studies in 2015. These studies collectively suggest that PRP and PRF are beneficial in reducing postoperative complications and enhancing the repair of soft tissue damaged during wisdom tooth surgery.

Conclusion: PRP and PRF effectively reduce postoperative complications and promote the restoration of soft tissues damaged by wisdom tooth extraction. Therefore, in cases of soft tissue damage and periodontal conditions in patients undergoing wisdom tooth surgery, PRP and PRF can be a valuable treatment option for tissue repair.

Keywords: Impacted wisdom teeth; Soft tissue; Hard tissue; Repair; Platelet-rich plasma (PRP); Platelet-rich fibrin (PRF).

Please cite this Article as:

Bolandparva F, Malekhosseini N, Asgarzade A. The Role of Platelet-Rich Products in Soft Tissue Repair after Extraction of Wisdom Teeth: A Review. J Craniomaxillofac Res 2025; 12(1): 1-6. DOI:



Copyright © 2025 Tehran University of Medical Sciences.

This work is licensed under a Creative Commons Attribution-NonCommercial 4.0 International license (<https://creativecommons.org/licenses/by-nc/4.0/>). Non-commercial uses of the work are permitted, provided the original work is properly cited.

Introduction

The wisdom tooth refers to the last tooth to erupt in each jaw arch, and if it is not placed in the dental arch at the expected time, it is called an impacted tooth. The most common impacted teeth are the upper and lower third molars. As a general rule, all impacted teeth should be extracted unless there are specific contraindications to extraction [1]. Surgical extraction of the third molar is one of the most common clinical procedures performed by surgeons [2]. This surgery is associated with the possibility of several complications after the operation, including pain, trismus, swelling, and dry socket (alveolar osteitis, AO) [3]. The prevalence of wisdom teeth in the lower jaw is 17.5%, and in the upper jaw, it is 21.9% [4]. Not removing these teeth from the jaw can lead to complications, including resorption of the roots of adjacent teeth, damage to the periodontal tissues, decay in adjacent teeth, and the development of cysts and tumors. Considering these potential issues, it appears that the most effective treatment method in such cases is to extract the teeth [5].

Socket repair of an avulsed tooth is a coordinated process of biochemical, physiological, cellular, and molecular reactions involving various cell types, growth factors, hormones, cytokines, and other proteins. This process aims to regenerate tissue and restore function after surgery. Following tissue damage, after hemostasis and cessation of bleeding, the repair process begins, which includes three interwoven stages: inflammation, fibroblastic activity, and regeneration. These stages overlap in terms of time [6]. Platelets are responsible for several vital activities, including blood coagulation, inflammation, antimicrobial defense, and wound healing. Although the initial focus of platelets is on coagulation, it is known that platelets carry growth factors, immune system messengers, enzymes, and other biologically active compounds involved in various aspects of tissue repair [7].

Platelets are involved in both hemostasis and the initiation of the wound healing process. These two processes are inseparable, as the tissue repair process begins concurrently with the completion of hemostasis [8]. Platelet-Rich Plasma (PRP) is obtained through the centrifugation of the patient's blood. It contains a physiological concentration of platelets that release growth factors. PRP is a biological approach to improve healing by releasing growth factors directly into the wound. In surgery, PRP has yielded beneficial results, including reduced bleeding, improved soft tissue repair and bone

regeneration, and numerous applications in treating musculoskeletal injuries. Animal and laboratory studies have demonstrated that PRP creates a local environment rich in growth factors and other cytokines for tissue regeneration, positively affecting the migration and proliferation of all cell types. Recently, expanding research on the effects of PRP regeneration on various tissues, such as bone, cartilage, tendon, and muscle, especially in traumatic injuries, has led to its increased use in plastic surgeries [9].

Studies conducted on humans using PRP in oral and dental surgeries, such as tooth extraction, periodontal surgery, and implant surgery, have shown promising results [10,11]. Research on the effects of PRP in improving wound healing in oral and maxillofacial surgery has yielded varied results, with most studies focusing on graft healing and dental implants [12]. PRP, which contains growth factors, can help prevent the development of periodontal defects caused by third molar extraction and also aid in their repair. During tooth extraction, bone formation typically takes approximately 16 weeks; however, there may be insufficient bone volume for reconstruction in the long term. PRP has been developed as an effective method to improve bone formation [13].

Platelet-Rich Fibrin (PRF) is prepared by a one-step centrifugation without applying any anticoagulants. PRF consists of platelets, leukocytes, and their subgroups embedded in a fibrin matrix with plasma proteins. The first protocol of PRF was called leukocyte-rich platelet-rich fibrin (L-PRF), mainly because it contains more leukocytes compared to the first-generation blood concentrates, PRP. PRF is the most extensively studied type of platelet concentrate for accelerating wound healing, particularly in promoting soft tissue migration. PRF can reduce inflammation of the periodontal tissues, preserve the alveolar site, and repair alveolar bone defects, thereby enhancing bone regeneration. L-PRF promotes rapid neo-angiogenesis and stimulates bone regeneration by releasing growth factors and providing good clot stability [14,15]. Considering the complications that can occur after wisdom tooth surgery, which significantly affect the patient's quality of life, as well as the effectiveness and efficiency of PRP and PRF, the purpose of this study is to review the role of PRP and PRF in soft tissue repair following wisdom tooth extraction from 2015 to 2023.

Materials and Methods

This research examined the results obtained from existing articles on the subject of a review of the role of

PRP and PRF in soft tissue repair after the extraction of impacted wisdom teeth. In this way, the articles available in the Scopus, PubMed, Elsevier and SID websites with similar topics in the 8 years from 2015 to 2023 were selected and reviewed based on the entry and exit criteria, and the articles were received by searching for wisdom teeth, soft tissue, repair, PRP and PRF keywords. The inclusion criteria for the study were: studies written in English, clinical studies, case-control studies, and review studies, conducted between 2015 and 2023, and articles with full-text availability. Exclusion criteria for the study were: systematic review studies and studies not indexed in the specified scientific databases. After searching the articles and reading the title and abstract, the necessary information (author, year of publication, type of study, number of patients, investigated parameter, type of treatment, conclusion) was obtained from the reviewed articles and the results were interpreted and compared.

Results

Based on the search strategy and inclusion criteria, 19 studies with the most similarity to the subject were identified, out of which no study had been conducted in 2023. The studies reviewed in this period were meta-analyses with low similarity to the subject of the study. Also, 3 studies in 2022, 4 studies in 2021, 3 studies in 2020, 1 study in 2019, 1 study in 2018, 1 study in 2017, 2 studies in 2016, and 4 studies in 2015 were reviewed. They had discussed the issue (Table 1).

Discussion

In most cases, wisdom tooth surgery creates problems and complications for the patient, which creates the need for repeated visits to treat and solve these problems. Common complications after wisdom tooth surgery include swelling, pain, bleeding, dry mouth, damage to soft tissue and periodontal structure and jaw bone. In studies conducted on human samples, the effect of PRP and PRF in oral and dental surgeries such as tooth extraction, periodontal surgeries and implant surgeries was promising. However, although many studies have investigated the effect of using PRP and PRF in various treatments and dental practices, but few studies have investigated the effect of PRP and PRF on soft tissue repair after the extraction of wisdom teeth. Collecting the latest achievements in this regard, reviewed the role of PRP and PRF in soft tissue repair after wisdom tooth extraction. Platelet-derived growth factors in PRP and PRF, as mitogens, play a key role in angiogenesis and tissue repair. Since the introduction

of PRP for maxillofacial use in 1997 by Withman, no single protocol has demonstrated significant, reproducible results. In a recent meta-analysis, limited evidence was found regarding the effects of PRP on bone repair, and the scientific evidence for PRP improving third molar cavities was also weak; therefore, further research is needed to fully identify its indications and efficacy in patients [16]. However, another study found improvement in bone and soft tissue after tooth extraction in patients [17]. In this review, 19 studies were identified that investigated the effect of PRP and PRF on soft tissue repair following third molar surgery. Also, other complications, including swelling, trismus, pocket depth and bone formation, were among the cases that were investigated in the studies. In most studies, there is a consensus that PRP and PRF resulted in soft tissue repair [18-31]. However, in some studies, although the soft tissue repair was more pronounced in the PRP and PRF group, no significant difference was observed between the two groups receiving PRP and PRF injections and the control group. In a study by Sargaiyan et al., most patients experience healing without the need for intervention, and PRP can only be effective in accelerating the healing of damaged soft tissue. But regarding pain caused by surgery, a positive effect of PRP injection was observed. Perhaps the reason for the lack of significant difference between the two investigated groups in the study of Sargaiyan et al. is due to the high periodontal health in the investigated patients in both groups and also the genetics of the patients is an effective factor in the restoration of soft tissue damaged by wisdom tooth surgery [32].

Another reason for the low effectiveness of PRP in soft tissue repair in some studies could be due to the nature of PRP and its effectiveness. In several studies, the effect of using PRP and PRF in soft tissue repair has been investigated [21,25,27,29,33,34]. They expressed the therapeutic effect of PRF in terms of soft tissue repair, pain, swelling, bleeding and bone formation far more than PRP. Osagie et al. stated in their study that PRF was more effective than PRP in reducing preoperative and postoperative swelling, the degree of mouth opening, and postoperative pain. In addition, this study demonstrated that platelet concentrate had a positive effect on reducing the inflammatory sequelae of latent brain surgery [33]. Bhujbal and colleagues also stated in their study that there was a significant improvement in soft tissue wound healing and an increase in bone density in the PRF site compared to PRP and a significant reduction in swelling on the first and third day in the PRF site compared to the PRP site.

Although postoperative pain scores were lower at the PRF site, it was not statistically significant between the two groups [21]. Doiphode et al. also observed a better therapeutic effect in the PRF group than PRP and control [27]. Dutta et al. also stated in their study, Yelamali et al. third molars were significantly better compared to PRP [28,29]. One of the reasons for the better effect of PRF compared to PRP may be due to the differences in their structure and production methods. The reason that the effect of PRF was greater than PRP may be due to the nature of PRF and the concentration of platelets in it compared to PRP. In comparisons made for the effect of PRP and PRF on tissue regeneration, it was stated that the effect of PRF on tissue regeneration is far greater than that of PRP [35,36]. It can also be attributed to the presence of leukocytes in PRF, as noted in several studies [35,37,38]. As Yelamali et al. stated in their research, the better effect of PRF compared to PRP may be due to the simpler preparation protocols of PRF compared to PRP and the ability of PRF to release growth factors in a controlled manner [29].

However, in some studies, no significant difference was observed between soft tissue repair in two PRF and PRP groups [34,27]. Verma et al. stated in their research that platelet-rich fibrin and platelet-rich plasma are both effective in reducing postoperative pain, swelling, and trismus, and no significant difference was observed. Both studied substances seem to have almost equal potential in bone and soft tissue healing [34]. Also, Dutta et al. in their study, which compared platelet-rich plasma (PRP), platelet-rich fibrin (PRF), and hydroxyapatite (HA) in soft tissue healing after wisdom tooth surgery, stated that PRP and PRF are better graft materials than HA in terms of pain, swelling, dry socket, and soft tissue healing. However, in comparison to other graft materials, bone regeneration is rapidly induced by HA [36].

Nevertheless, a more precise study with a larger number of clinical cases is necessary to determine the definitive effectiveness of graft materials. The genetic makeup of individuals can significantly impact the healing of soft tissues following surgery, which could explain the variability in study results. Additionally, factors such as the technician's skill and the extent of the injury are not measurable and were not addressed in previous studies. However, the therapeutic effect of PRP and PRF may vary in patients with different surgical conditions and levels of soft tissue damage. Therefore, based on the results of a single study, it is not possible to conclusively confirm or refute the effect of

PRP on soft tissue healing. A comprehensive assessment of study results is necessary to make an informed decision regarding the use or non-use of PRP. Out of 19 studies reviewed, 18 studies indicated that PRP and PRF are effective in soft tissue healing and reducing other complications following wisdom tooth surgery. Only one study did not find a significant difference between the two groups under study. Still, even in this study, the soft tissue healing effect in the PRP group was greater than in the control group. Thus, based on a consensus, PRP and PRF can be considered effective in soft tissue healing. It should also be noted that the use of PRP and PRF should be based on the surgeon's discretion, careful assessment of the extent of the injury, the individual's periodontal disease history, and the type of surgery being performed. Hence, the necessity of using PRP and PRF in all patients is not obligatory, but its benefits should be utilized when needed.

Conclusion

PRP leads to reduction of complications after surgery and restoration of soft tissue damaged by wisdom tooth surgery; therefore, in case of damage to soft tissue and periodontal condition in patients undergoing wisdom tooth surgery, this method can be used to restore soft tissue.

Conflict of Interest

There is no conflict of interest to declare.

References

- [1] Hupp JR, Tucker MR, Ellis E. Contemporary otal and maxillofacial surgery. china, Mosy, 2008, pp, 149-184.
- [2] Dodson TB, Susarla SM. Impacted wisdom teeth. *BMJ Clin Evid* 2014; 2014:1302.
- [3] Candotto V, Oberti L, Gabrione F, Scarano A, Rossi D, Romano M. Complication in third molar extractions. *J Biol Regul Homeost Agents* 2019; 33(3 Suppl 1):169-72.
- [4] Rakhshan V. Common risk factors for postoperative pain following the extraction of wisdom teeth. *Journal of the Korean Association of Oral and Maxillofacial Surgeons*. 2015 Apr 1; 41(2):59-65.
- [5] Miloro M. Petersons Principles of Oral and Maxillofacial Surgery. 2th ed. Canada: BC Decker In; 2004. P. 140-2.

- [6] Hupp JR, Tucker MR, Ellis E. Contemporary oral and maxillofacial surgery. 7th ed. St. Louis, Missouri: Elsevier; 2018. p. 189, 190.
- [7] Leslie M. Cell biology. Beyond clotting: the powers of platelets. *Science* 2010; 328(5978): 562-4.
- [8] Scimeca CL, Bharara M, Fisher TK, Kimbriel H, Armstrong DG. Novel use of platelet-rich plasma to augment curative diabetic foot surgery. *J Diabetes Sci Technol* 2010; 4(5): 1121-6.
- [9] Simon D, Manuel S, Geetha V, Naik BR. Potential for osseous regeneration of platelet-rich plasma--a comparative study in mandibular third molar sockets. *Indian J Dent Res* 2004; 15(4): 133-6.
- [10] Albanese A, Licata ME, Polizzi B, Campisi G. Platelet-rich plasma (PRP) in dental and oral surgery: from the wound healing to bone regeneration. *Immun Ageing* 2013; 10(1): 23.
- [11] Marco Antonello Gd, Torres do Couto R, Giongo CC, Corrêa MB, Chagas Júnior OL, et al. Evaluation of the effects of the use of platelet-rich plasma (PRP) on alveolar bone repair following extraction of impacted third molars: prospective study. *J Craniomaxillofac Surg* 2013; 41(4): e70-5.
- [12] Ogundipe OK, Ugboko VI, Owotade FJ. Can autologous platelet-rich plasma gel enhance healing after surgical extraction of mandibular third molars? *J Oral Maxillofac Surg* 2011; 69(9): 2305-10.
- [13] Rutkowski JL, Johnson DA, Radio NM, Fennell JW. Platelet rich plasma to facilitate wound healing following tooth extraction. *J Oral Implantol* 2010; 36(1): 11-23.
- [14] Al-Maawi S, Becker K, Schwarz F, Sader R, Ghanaati S. Efficacy of platelet-rich fibrin in promoting the healing of extraction sockets: a systematic review. *International journal of implant dentistry*. 2021 Dec; 7:1-27.
- [15] Alrayyes Y, Al-Jasser R. Regenerative potential of platelet Rich Fibrin (PRF) in Socket Preservation in comparison with Conventional Treatment modalities: a systematic review and Meta-analysis. *Tissue Engineering and Regenerative Medicine*. 2022 Jun; 19(3):463-75.
- [16] Agonas P, Schiavo J.H., Avila-Ortiz G., Palaiologou A., Katsaros T. Plasma rich in growth factors (PRGF) in intraoral bone grafting procedures: a systematic review. *J Cran-Maxillo-Fac Surg*. 2019; 47(3):443-453.
- p17] Barona-Dorado C., Gonzalez-Regueiro I., Martin-Ares M., Arias-Irimia O., Martinez-Gonzalez J.M. Efficacy of platelet-rich plasma applied to post-extraction retained lower third molar alveoli. A systematic review. *Med Oral, Patol Oral Cirugía Bucal*. 2014; 19(2):142-148.
- [18] Rengarajoo J, Ngeow WC, Ibrahim NB. The effects of lyophilised platelet-rich plasma in third molar extraction sockets and its surrounding tissues. *Journal of Taibah University Medical Sciences*. 2022 Apr 1; 17(2):289-96.
- [19] Ahmed I, Hussain F, Naz S, Hassan SG, Channar KA, Shams S. Role Of Platelet Rich Plasma On Healing Following Surgical Extraction Of Teeth. *Journal of Peoples University of Medical & Health Sciences Nawabshah. (JPUMHS)*. 2021 Mar 31; 11(1):118-22.
- [20] Al-Noaman AS. Platelet-rich plasma in oral and dental surgery: A review. *Medical Journal of Babylon*. 2021 Apr 1; 18(2):59-65.
- [21] Bhujbal R, Veerabhadrappe SK, Yadav S, Chappi M, Patil V. Evaluation of platelet-rich fibrin and platelet-rich plasma in impacted mandibular third molar extraction socket healing and bone regeneration: A split-mouth comparative study. *European Journal of General Dentistry*. 2020 May; 9(02):96-102.
- [22] Aftab A, Joshi UK, Patil SK, Hussain E, Bhatnagar S. Efficacy of autologous platelet rich plasma gel in soft and hard tissue healing after surgical extraction of impacted mandibular third molar-A prospective study. *Journal of Oral and Maxillofacial Surgery, Medicine, and Pathology*. 2020 Jul 1; 32(4):241-6.
- [23] Jonathan R. The effect of lyophilized platelet rich plasma in a third molar extraction socket and its surrounding tissues/Jonathan Rengarajoo (Doctoral dissertation, Universiti Malaya).
- [24] Bhujbal R, Malik NA, Kumar N, Suresh KV, Parkar MI, Jeevan MB. Comparative evaluation of platelet rich plasma in socket healing and bone regeneration after surgical removal of impacted mandibular third molars. *Journal of Dental Research, Dental Clinics, Dental Prospects*. 2018; 12(3):153.
- [25] Gandevivala A, Sangle A, Shah D, Tejani A,

- Sayed A, Khutwad G, Patel AA. Autologous platelet-rich plasma after third molar surgery. *Annals of Maxillofacial Surgery*. 2017 Jul 1; 7(2):245-9.
- [26] Doiphode AM, Hegde P, Mahindra U, Kumar SS, Tenglikar PD, Tripathi V. Evaluation of the efficacy of platelet-rich plasma and platelet-rich fibrin in alveolar defects after removal of impacted bilateral mandibular third molars. *Journal of International Society of Preventive and Community Dentistry*. 2016 Apr 1; 6(Suppl 1):S47-52.
- [27] Dutta SR, Passi D, Singh P, Sharma S, Singh M, Srivastava D. A randomized comparative prospective study of platelet-rich plasma, platelet-rich fibrin, and hydroxyapatite as a graft material for mandibular third molar extraction socket healing. *National journal of maxillofacial surgery*. 2016 Jan 1; 7(1):45-51.
- [28] Dutta SR, Singh P, Passi D, Patter P. Mandibular third molar extraction wound healing with and without platelet rich plasma: a comparative prospective study. *Journal of maxillofacial and oral surgery*. 2015; 14:808-15.
- [29] Yelamali T, Saikrishna D. Role of platelet rich fibrin and platelet rich plasma in wound healing of extracted third molar sockets: a comparative study. *Journal of maxillofacial and oral surgery*. 2015; 14:410-6.
- [30] Gawai KT, Sobhana CR. Clinical evaluation of use of platelet rich plasma in bone healing. *Journal of maxillofacial and oral surgery*. 2015; 14:67-80.
- [31] Nathani DB, Sequeira J, Rao BS. Comparison of platelet rich plasma and synthetic graft material for bone regeneration after third molar extraction. *Annals of maxillofacial surgery*. 2015 Jul 1; 5(2):213-8.
- [32] Sargaiyan V, Manas A, Kumar HH, Saravanan MS, Ghiaz K, Deepalakshmi S. The role of prp in third molar extraction wounds: A clinical study. *Journal of Pharmacy and Bioallied Sciences*. 2022 Jul 1; 14(Suppl 1): S855-8.
- [33] Osagie O, Saheeb BD, Egbor EP. Evaluation of the Efficacy of Platelet-Rich Plasma versus Platelet-Rich Fibrin in Alleviating Postoperative Inflammatory Morbidities after Lower Third Molar Surgery: A Double-Blind Randomized Study. *West African Journal of Medicine*. 2022 Apr 1; 39(4):343-9.
- [34] Verma VK, Sublok K, Kushwaha AK, Patra D, Khare A, Singh J. A prospective study to evaluate post operative clinical parameters of platelet rich fibrin and platelet rich plasma in third molar extraction sockets: An original research study. *Journal of Advanced Medical and Dental Sciences Research*. 2021; 9(1):45-50.
- [35] Giannini, S., et al. Comparison between PRP, PRGF and PRF: lights and shadows in three similar but different protocols. *Eur Rev Med Pharmacol Sci*, 2015, 19.6: 927-30.
- [36] Dohan Ehrenfest DM, Del Corso M, Diss A, Mouhyi J, Charrier JB. Three-dimensional architecture and cell composition of a Choukroun's platelet-rich fibrin clot and membrane. *Journal of periodontology*. 2010; 42(55):62.
- [37] Dohan DM, Choukroun J, Diss A, Dohan SL, Dohan AJ, Mouhyi J, et al. Platelet-rich fibrin (PRF): a second-generation platelet concentrate. Part III: leucocyte activation: a new feature for platelet concentrates? *Oral surgery, oral medicine, oral pathology, oral radiology, and endodontics*. 6112; 010(3): e20-2.
- [38] Hoaglin DR, Lines GK. Prevention of localized osteitis in mandibular third-molar sites using platelet-rich fibrin. *International journal of dentistry*. 6103; 61033062301.
- [39] Hanif M, Sheikh MA. Efficacy of platelet rich plasma (PRP) on mouth opening and pain after surgical extraction of mandibular third molars. *Journal of Oral Medicine and Oral Surgery*. 2021; 27(1):9.
- [40] Kumar A, Singh G, Gaur A, Kumar S, Numan M, Jacob G, Kiran K. Role of platelet-rich plasma in the healing of impacted third molar socket: a comparative study on central India population. *J Contemp Dent Pract*. 2020 Sep 1; 21(9):986-91.