



Association between the serotonin- transporter- linked polymorphic region (5-HTTLPR) and pemphigous disease in Iranian patients

Shamsolmolok Najafi ¹, Nafiseh Esmaili ², Mansour Heidari ³, Farahnaz Ghassemi ⁴, Meisam Mahmoudi ⁵, Mahsa Mohammadzadeh ⁵, Abdolreza Mohamadnia ⁶, Naghmeh Bahrami ^{7*}

1. Oral Medicine department, Dental School, Tehran University of Medical Sciences, Tehran, Iran.

2. Skin & hair department, Razi hospital, Tehran University of Medical Sciences, Tehran, Iran.

3. Eye Research Center, Farabi Eye Hospital, Tehran University of Medical Sciences, Tehran, Iran.

4. Cranio-maxillofacial Research center, Tehran University of Medical Sciences, Tehran, Iran.

5. Dentist

6. Virology Research Center, National Research Institute of Tuberculosis and Lung Diseases (NRITLD), Shahid Beheshti University of Medical Sciences, Tehran, Iran. Biotechnology Department, advanced technologies in medicine, Shahid Beheshti University of Medical Sciences, Tehran, Iran.

7. Oral and Maxillofacial Surgery Department, School of Dentistry, Tehran University of Medical Sciences, Tehran, Iran. Cranio-maxillofacial Research center, Tehran University of Medical Sciences, Tehran, Iran, Iranian Tissue Bank & Research Center, Tehran University of Medical Sciences, Tehran, Iran.

ARTICLE INFO

Article Type:
Original Article

Received: 11 Jan 2016

Revised: 26 Feb 2016

Accepted: 29 Mar 2016

*Corresponding author:

Naghmeh Bahrami

Oral and Maxillofacial Surgery Department, School of Dentistry, Tehran University of Medical Sciences, North Amirabad St, Tehran, Iran

Tel: +98-21-84902473

Fax: +98-21-84902473

Email: NaghmehBahrami@Gmail.com

ABSTRACT

Introduction: Pemphigus vulgaris (PV) is a relatively rare autoimmune disease characterized by blistering of the skin and mucosa. The main objective of this study was to investigate the possible association between the 5-HTTLPR polymorphism and PV in Iranian patients. For this, 112 PV and 100 controls were enrolled in this study.

Materials and Methods: In this case-control study, Genomic DNA was extracted from whole blood and genotyping of all participants for the 5-HTTLPR polymorphism was carried out using the polymerase chain reaction (PCR) technique. The genotypes were grouped into three classes: homozygous for the short allele (SS), heterozygous for the short and long allele (LS) and homozygous for the long allele (LL).

Results: Our results showed no significant association between the 5-HTTLPR polymorphism genotypes, LL, SS and LS in PV patients compared to controls. Also, our finding did not reveal any evidence of an association between this disease and the allele frequency of S and L. Taken together, our findings suggested that the 5-HTTLPR polymorphism is unlikely to be a factor contributing to the risk of developing pemphigus vulgaris.

Conclusion: It can be concluded that although 5-HTTLPR polymorphism seems to be associated with some of auto-immune and stress-related disease.

Key words: Iranian Population, PCR, Pemphigus vulgaris, Polymorphism, Risk Factor.

Introduction

Pemphigus vulgaris (PV) as a rare and autoimmune disease causes very serious blisters on skin as well as inside the mouth, nose and throat [1]. So far, several types of this disorder have been reported such as mucosal pemphigus vulgaris, muco-cutaneous pemphigus

vulgaris, paraneoplastic pemphigus, drug-induced pemphigus, pemphigus foliaceus, and IgA pemphigus [2]. The most prevalent variant is pemphigus vulgaris (PV) [3], which its annual incidence is about 1/100000 in Iran, with a mean age of onset of 42 years, a female to male ratio of

1.5/1 and a death rate of about 6% [4].

In PV, autoantibodies are produced against such desmosome components of epithelium as cadherin and desmoglein (Dsg), especially Dsg3 [2]. These glyco-protein molecules are found in a membrane in desmosome and strengthen intercellular connection. The lack of these connections due to the antibody-antigen reactions leads to weakness of intercellular connection, and finally separation of the cells, which in turn results in desquamation and blister formation [5]. While Dsg1 and Dsg3 both are expressed in skin, oral epithelium expresses predominantly only Dsg3 [6]. Thus, the first manifestation of the disease is mucosal involvement among 74% of the cases [7]. Moreover, up to 90% of the patients present oral lesion as the first sign of the disease development [5].

Previous studies have reported that various etiological factors such as, cosmetic materials [15], estrogen hormone [16] diet [8], such drugs as thiol [9] and non-thiol drugs [10], phenol drugs [11], rifampicin [12], diclofenac [13], and other angiotensin-converting-enzyme inhibitors [14] contribute to the pathogenesis of PV. In addition, it is well known that psychological stress may exacerbate skin diseases [17,18].

There is a body of evidences showing that psychological status is affected by serotonergic mechanisms especially serotonin-transporter mechanisms (5-HTT) [19]. Since 5-HTT regulates serotonergic neurotransmission, polymorphism in this gene is associated with anxiety-related behaviours [20, 21]. 5-HTT (SLC6A4) gene is located on chromosome 14q12 and contains promoter and 14 Kb31 spanning exons [22, 23]. Two polymorphic regions, a 44-bp insertion/deletion polymorphism within the promoter region (5-HTTLPR) and a 17-bp variable number of tandem repeat polymorphism (VNTR) in intron 2 [22, 24] in SLC6A4 have been demonstrated. These allelic forms, the long (L) and the short (S) variants are found in the first polymorphic region (5-HTTLPR) [22]. It has been reported that the two alleles differently regulate SLC6A4: S allele reduces the transcriptional efficiency of the 5-HTT, and consequently decreases uptake of serotonin [20], whereas L allele has the opposite effect of S allele.

In spite of the fact that several studies have been conducted on the association between various types of polymorphisms with PV, to our knowledge, none of them has investigated the association of 5-HTT polymorphism with PV.

The aim of the present study was to investigate 5-HTTLPR polymorphism in 5-HTT gene in the Iranian patients with PV.

Materials and Methods

In this case-control study, we investigated a total of 212 Iranian subjects comprising 112 PV cases and 100 healthy controls. All patients were seen at Skin Disease Ward, Razi Hospital, Tehran, Iran. The participants gave their written informed consent, and the study was approved by the TUMS ethics review board as part of the reviewing process of TUMS research projects. The study was conducted according to the principles of the Helsinki Declaration.

All individuals underwent a complete clinical examination. The diagnosis of PV was made by a dermatologist, further confirmed by biopsy. The size of the lesion was measured, and the duration of the disease as reported by the patient was recorded.

Genotyping

Five ml blood sample of all participants were collected, and DNA genome was extracted using a Pure Nucleic Acid Isolation Kit (Roche Applied Science, Mannheim, Germany). Genotyping of all individuals for the 5-HTTLPR polymorphisms were conducted using the PCR technique. PCR amplification was performed using specific primers: 5HTTF: 5'-TGAATGCCAGCACCTAACCC-3'; 5HTTR: 5'-TTCTGGT-GCCACCTAGACGC-3') 1U Taq DNA polymerase (Roche, Mannheim, Germany), 10 pM of each primer, 200 µM of each dNTPs, 0.67 µl of 50 mM MgCl₂, 60 ng DNA and 2.5 µl of 10X PCR buffer in 25 µl PCR reactions. The PCR was performed based on the following conditions: initial denaturation at 94°C for 5 min; followed by 35 cycles including denaturation at 94°C for 35 s, annealing at 58°C for 35 s, and extension at 72°C for 1.5 min; and a final extension at 72°C for 5 min. In order to identify S and L alleles in 5-HTTLPR polymorphism, PCR products (400bp and 450 bp, respectively) were visualized by agarose gel electrophoresis. All the subjects were divided into three genotype groups: homozygote for S alleles (SS), heterozygote for S and L alleles (LS), and homozygote for L alleles (LL).

Statistical analyses

SPSS version 11.2 served for statistical analyses. Association between 5-HTTLPR and PV was tested by Fisher's exact test at 0.05 significance level.

Results

The mean duration of illness among PV patients was 7.9 years (95%CI: 6.8-8.9) and the mean size of their lesions was 10 mm (95% CI: 9.6-12.1). The principal characteristics of the subjects in case and control groups are shown in the Table 1.

Genotypes in our study were identified using PCR . The most frequent genotype among both groups was SL (about 60%), followed by LL (20%), and SS (less than 20%). The distribution of the genotypes between the two groups showed no statistically significant difference ($P=.791$). The frequency of three genotypes of 5-HTTLPR in case and control groups is shown in Table 2.

No statistically significant difference was found between the two groups in this regard ($P=.553$). Table 3 shows the frequency of L and S alleles of 5-HTTLPR in case and control groups.

Discussion

In the present study identified the genotypes of 112 patients with PV and 100 healthy subjects (SS homozygote, LS heterozygote, and LL homozygote) using specific primers of 5-HTTLPR in a PCR technique. According to the results no statistically significant differences existed between the case and control groups regarding the frequency of the three genotypes, and also the frequency of S and L alleles. These findings suggest neither that frequency of neither genotypes nor alleles of 5-HTTLPR could be a risk factor for PV.

Association between serotonin transporter promoter polymorphism (5-HTTLPR) and autoimmune diseases has been frequently reported, though it has been a matter of controversy in the last decade. The main problem in genetic studies on PV as a multi-factorial disease is probably contribution of various factors such as psychological stress to disease development. Moreover, clinical diagnoses may be multi-factorial constructs that combine elements with distinct genetic influences.

A study on stressful life events, 5-HTTLPR, and major depression found no evidence of either genetic support for the disease, or interaction between genetic and environmental factors [25]. Another study could not relate signs of mental disorders and stress factors to 5-HTTLPR among children exposed to urban violence [26]. On the other hand many studies have related anxiety and major depression disorders to 5-HTTLPR genotype [27-29]. For example, a meta-analysis

by Srijan Sen et al. reported significant association between 5-HTTLPR and personality-related anxiety [30]. A systematic review in US investigated association of polymorphism in 5-HTTLPR with anxiety-related behaviours. Meta-analysis was done on 23 studies and the results indicated a strong relationship between 5-HTTLPR polymorphism and neuroticism-related behaviours [31].

Victoria et al studied serotonin transporter gene polymorphism among 69 patients with recurrent aphthous stomatitis (RAS) and 70 healthy subjects and found remarkable increase in frequency of SS genotype, S alleles, and 5-HTTLPR polymorphism among RAS patients compared to healthy controls [32].

In conclusion, our data suggest that the 5-HTTLPR polymorphism is not a genetic risk for the development of PV within the Iranian population. Although the 5-HTTLPR variant (L) is associated with an increased expression of serotonin levels in the disease with psychological stress, other molecular genetics could be implicated in the initiation, progression and development of PV.

Conclusion

It can be concluded that although 5-HTTLPR polymorphism seems to be associated with some of auto-immune and stress-related disease, we could not find such an association for PV among Iranian population. Future studies can be done among various ethnic or age groups, with larger sample sizes.

Table 1. Distribution of background characteristics of patients with pemphigus vulgaris (case group, n=112) and healthy subjects (control group, n=100).

Characteristics	PV (n=112)	Control (n=100)
Age (year): Mean (95% CI)	41.5 (18 - 65)	38 (20 - 56)
Minimum	18	20
Maximum	65	56
Lesion Size: Mean (95% CI) (cm ²)		
Gender: Female , Male		

Table 2. The frequency of three genotypes of serotonin-transporter-linked polymorphic region (5-HTTLPR) in patients with pemphigus vulgaris (case group), and healthy subjects (control group).

	LL n (%)	SL n (%)	SS n (%)
Case group (n=112)	25 (23.3)	69 (61.6)	18 (16.0)
Control group (n=100)	26 (26.0)	60 (60.0)	14 (14.0)
Total	51	129	32

Table 3. The frequency of L and S alleles of serotonin-transporter-linked polymorphic region (5-HTTLPR) in case (patients with pemphigus vulgaris) and control (healthy subjects) groups.

	S allele	L allele
Case group (n=112)	105 (46.9%)	119 (53.1%)
Control group (n=100)	88 (44.0%)	112 (56.0%)

References

- [1] Veldman C, Feliciani C. Pemphigus: A Complex T Cell-dependent Autoimmune Disorder Leading to Acantholysis. *Clinic Rev Allerg Immunol*. 2008; 34: 313-320.
- [2] Scully C, Challacombe SJ. Pemphigus vulgaris: update on etiology, oral manifestations, and management. *Oral Manifestations, and Management. CROBM*. 2002; 13(5): 397-408.
- [3] Thorat MS, Raju A, R. Pradeep A. Pemphigus vulgaris: effects on periodontal health. *Journal of Oral Science*. 2010; 52(3): 449-454.
- [4] Chams-Davatchi C, Valikhani M, Daneshpazhooh M, Esmaili N, Hallaj Z, Barzegari M, Akhiani M, Ghodsi Z, Mortazavi H, Naraghi Z. Pemphigus: Analysis of 1219 cases. *International Journal of Dermatology*. 2014; 53:531-553.
- [5] Burket LAW, Greenberg MS, Glick M. *Vesicular and Bullous Lesion. Burket's Oral Medicine*. 11th ed. United states: Bc Decker. 2003; 211-232.
- [6] Shirakata Y, Amagai M, Hanakawa Y, Nishikawa T, Hashimoto K. Lack of mucosal involvement in pemphigus foliaceus may be due to low expression of desmoglein 1. *Journal of Investigative Dermatology*. 1998; 110: 76-78.
- [7] Asilian A, Yoosefi A, Faghini G. Pemphigus vulgaris in Iran: epidemiology and clinical profile. *SKINmed: Dermatology for the Clinician*. 2006 March/April; 5(2): 69-71.
- [8] Brenner S, Ruocco V, Bialy-Golan A, et al. Drug-induced pemphigus. *Clinics in dermatology*. 2011; 39: 313-393.
- [9] Shapiro M, Jimenez S, Werth VP. Pemphigus vulgaris induced by D-penicillamine therapy in a pa-

- tient with systemic sclerosis. *J Am Acad Dermatol*. 2000 Feb; 42:297-9
- [10] Wolf R, Brenner S. An active amide group in the molecule of drugs that induce pemphigus: a casual or causal relationship?. *Dermatology* .1994;189:1-4.
 - [11] Goldberg I, Kashman Y, Brenner S. The induction of pemphigus by phenol drugs. *International Journal of Dermatology*. 1999 December; 38(12): 888-892.
 - [12] Gange R.W., Rhodes E.L., Edwards C.O., Powell M.E. A. Pemphigus induced by rifampicin. *British Journal of Dermatology*. October 1976; 95(4):445-448.
 - [13] Matz H, Bialy-Golan A, Brenner S. Diclofenac: a new trigger of pemphigus vulgaris?. *Dermatology* .1997; 195:48-49.
 - [14] Ong CS, Cook N, Lee S. Drug-related pemphigus and angiotensin converting enzyme inhibitors. *Australasian journal of dermatology*. 2000 November; 41(4):242-246.
 - [15] Bastuji-Garin S, Turki H, Mokhtar I, Nouira R, Fazaa B, Jomaa B, et al. Possible relation of Tunisian pemphigus with traditional cosmetics: A Multicenter case-control study. *American Journal of Epidemiology*. 2001; 155(3): 249-256.
 - [16] Brenner S, Tur E., Shapiro J, Ruocco V, D'Avino M, Ruocco E, et al. Pemphigus vulgaris: environmental factors. Occupational, behavioral, medical, and qualitative food frequency questionnaire. *International Journal of Dermatology*. 2001 September; 40(9): 562-569.
 - [17] Kimayi-Asadi A , Usman A. The role of psychological stress in skin disease. *Journal of Cutaneous Medicine and Surgery*. 2001; 4(2):140-145.
 - [18] Morell-Dubois S, Carpentier O, Cottencin O, Queyrel V, Hachulla E. Stressful life Events and Pemphigus. *Dermatology*. 2008; 216(2):104-108.
 - [19] Victoria JMN, Fátima CS, Flávio JP, Kalapothakis E, Gomez RS. Serotonin transporter gene polymorphism (5-HTTLPR) in patients with recurrent aphthous stomatitis. *Journal of Oral Pathology & Medicine*. 2005 September; 34(8): 494-497.
 - [20] Lesch KP, Bengel D, Heils A, et al. Association of anxiety-related traits with a polymorphism in the serotonin transporter gene regulatory region. *Science magazine*. 1996 November; 274: 1527-1531.
 - [21] Greenberg B.D, Li Q, Lucas FR, et al. Association between the serotonin transporter promoter polymorphism and personality traits in a primarily female populations sample. *American Journal Of Medical Genetics*. 2000 April; 96 (2): 202-216.
 - [22] Lesch K.P, Balling U, Gross J, et al. Organization of the human serotonin transporter gene. *Journal Of Neural Transmission*. 1994; 95: 157-162.
 - [23] Ramamoorthy S, Bauman AL, Moore KR, Han H, Yang-Feng T, Chang AS. Antidepressant and cocaine sensitive human serotonin transporter: localization molecular cloning, expression and chromosomal localization. *Proc Natl Acad Sci U S A*. 1993 Mar 15; 90 (6):2542-6.
 - [24] Heils A, Teufel A, Petri S, et al. Allelic variation of human serotonin transporter gene expression. *Journal Of Neurochemistry*. 1996 June; 66 (6): 2621-2624.
 - [25] Caspi A, Sugden A, E. Moffitt t, et al. Influence of Life Stress on Depression: Moderation by a Polymorphism in the 5-HTT Gene. *Science*. 2003 Jul 18; 301 (5631):386-9
 - [26] C. Cividanes G, F. Mello A, M. Sallum J, et al. Lack of association between the 5-HTTLPR and positive screening for mental disorders among children exposed to urban violence and maltreatment. *Revista Brasileira de Psiquiatria*. 2014; 36: 277-284.
 - [27] Kendler KS, Kuhn JW, Vittum J, Prescott CA, Riley B. The interaction of stressful life events and a serotonin transporter polymorphism in the prediction of episodes of major depression: A replication. *Arch Gen Psychiatry*. 2005; 62:529-35.
 - [28] Eley TC, Sugden K, Corsico A, Gregory AM, Sham P, McGuffin P, et al. Gene-environment interaction analysis of serotonin system markers with adolescent depression. *Molecular Psychiatry*. 2004; 9:908-915.
 - [29] Caspi A, Hariri AR, Holmes A, Uher R, Moffitt TE. Genetic sensitivity to the environment: the case of the serotonin transporter gene and its implications for studying complex diseases and traits. *Am Psychiatric Assoc*. 2010 Summer; 8(3):398-416.
 - [30] Srijan Sen, Margit Burmeister, Debashis Ghosh.

Meta-Analysis of the Association Between a Serotonin Transporter Promoter Polymorphism (5-HTTLPR) and Anxiety-Related Personality Traits. American Journal of Medical Genetics Part B: Neuropsychiatric Genetics. 2004 May; 127B (1): 85-89.

- [31] Natan A, Gillespie, John B. Whitfield, Ben Williams. The relationship between stressful life events, the serotonin transporter (5-HTTLPR) genotype and major depression. Psychological Medicine. 2005; 35(1):101-111.
- [32] Harman KE, Albert S, Black MM. Guidelines for the management of pemphigus vulgaris. British Journal of Dermatology. 2003; 149 (5):926-937.

Please cite this paper as:

Najafi Sh, Esmaili N, Heidari M, Ghassemi F, Mahmoudi M, Mohammadzadeh M, Mohamadnia A, Bahrami N; Association between the serotonin-transporter-linked polymorphic region (5-HTTLPR) and pemphigous disease in Iranian patients. J Craniomaxillofac Res 2016; 3(2): 185-190